

Toboa na kupondoponda kabisa mitebe na mapipa ya sheria za utopaji. Ustumiwe mitebe au mapipa ya dawa kwa kazi yoyote, na hasa kwa kuweka chakula, malisho ya mifugo au maji.

Usalama kwa mtumiaji

EPUKA kugusa na dawa kwenye mwili, macho, ama kwenye nguo. Usitute hewa yenye vumbi la dawa. Usiile, kunywa au kuvuta sigara wakati wa uchanganyaji. Nawa mikono na uso kabla ya kula kunywa au kuvuta sigara.

Vaa mavazi ya kinga: overoli, glavu zizozipitisha maji, na buti. Baada ya kazi, osha nguo na vifa vya kazi vilivyotumika. Osha mwili mzima na kicha badilisha nguo. Fua nguo iliyomwagika dawa kabla ya kutumia tena.

Mazingira na wanyama wa pori

Usitute mabaki ya dawa kwenye maji yatumikayo kwa au matumizi ya nyumbani na yezzo, vingine vya maji kama vile mabwazi, mifereji, mazwa, n.k.

Kuvuja na Kumwagika dawa.

Udongo uliomwagika dawa uchotwe viziti na utupile mahali salama mbali na vianzo au njia za maji. Kusanya dawa iliyomwagika kicha weka kwenye mifuko wa plastiki na kutupa sehemu iliyomwagika dawa kwa maji mengi.

HUDUMA YA KWANZA

Iliwauwa punzi: Unga wa ACTELIC SUPER waweza kusababisha hali ya kupiga chalya na kukotoa iwapo utapewa mwanya wa kuingilia njia za upumaji. Mitu mwenye kuhusha haraka lazima aingie sehemu atabakopajita hewa safi.

Nguzwi: Mwiili uliochafuwa na dawa uoshwe kwa maji na sabuni.

Machoni: Miruko machoni lochw kwa maji safi kwa muda wa angalau dakika 15. Muone daktari.

Ikiwa imecwa: Endapo ACTELIC SUPER imeme zwa usimlazimisho mgonjwa kutapika, haraka mpeleke mgonjwa pamoja na kibandiko hiki kwa daktari.

PELEKE MIERUHI KWA DAKTARI

Dalli za sumu: Hizi ni pamoja na kuchefuchefu, kukumba na kichwa, kutoona viziti, mbano wakifuani, ulgevu wa viungo na maumivu ya tumbo. Hali ikiwa mbaya zaidi hueda kukawa na tatizo kama kupumua, perigine itakuwa lazima kumsaidi kupumua.

Kinga: Atropine sulphate ACTELIC SUPER ni dawa iliyotengenezwa kwa misingi wa pyrethroid na ya fosforasi. Iwapo italithwa shaka kuwepo kwa sumu, dawa ya Atropine itolewe mara moja kwa kudungwa sindano.

na mshpa na mtu mwenye ujuzi kwa kipimo cha mg 2.5 kifuatiwa na mg 2 bila nusu saa. Nakati huo huo dawa ya Pralidoxime (p-25 au P-2m) itolewe. kwa kipimo cha gramu 1 kwa mililita 2.3 ya maji kwa sindano inayodungwa kwenye mshpa. Zaitolewe dawa za kutuliza. Ruda matumizi ya dawa kutegemea maendeleo ya matibabu.

Kukiwa na jambo la dharuru piga simu isiotozwa Nambari: 0800 720021 / 0800 730030 (Masa 24)

ILANI KWA MTUMIAJI

Dawa hii ni sharti itumike kulingana na maagizo yaliyo kwenye kibandiko. Ni hasa chini ya sharia za madawa ya kuangiza wadudu waharibifu kutumia au kuhifadhi dawa hii katika hali isyo salama.

THIBITISHO

Dhamana ya muuzaji ni maelezo yaliyo kwenye kibandiko ambayo ni lazima yafuatwe kikamilifu. Mfunuzi lazima akubali hatari zozote zinazoweza kutokana na matumizi ya dawa hii kwa binadamu au maji na lazima akubali dawa hii katika hali isyo.

Broad spectrum insecticidal dust for the control of stored maize and small grain cereals including control of larger grain borers.

Maelezo

Actellic Super Dust ni kiuadudu aina ya vumbi yenye pirimiphos-methyl iliyokwenye kundi la organophosphate, actellic Super Dust ina umalifu wa ngusio na sumu tumbo. Pia hubaki kwenye punje na kutoa hewa ya sumu.

Namba ya umalifu

Pirimiphos-methyl hupenya haraka kwenye mwili wa mndu na kuzua utendaji kazi wa acetylcholinesterase. Permethrin hutenda kazi katika mshpa ya niva kwa mndu anayehusika.

Matumizi kwa mazaio yaliyoko ghalani Actellic Super Dust hudhibiti wadudu wenji muhimu washambuliao mazaio ghalani kama vile wadudu walomo kwenye maji ya nondo, wadudukobe, wadudutumbo. Hudhibiti wadudu walogo na kukinga dhidi ya mashambizi mengine ya baada.

Actellic Super Dust hudanya kazi katika mazingira ya joto ya nchi za tropiki. Hubaki kwenye punje za nafaka kwa mizi kadhaa Actellic Super Dust yaweza kurunywa moja kwa moja kwenye nafaka na kwenye magunga.

JINSI YA KUTUMIA

Kwa matokeo bora nafaka zitilwe dawa pale zinapoenleka kutiwa ghalani. Uharibifu ukioleka na kushika sana si rahisi kuuzulu. Ingawa ni muhimu kueneza dawa kwenye nafaka zote, lakini kwa kiasi fulani ule uwezo wake wa kufanya kazi kwa msambao wa husaidia kueneza dawa. Mbebu za nafaka zitizothe dawa hukingwa kwa muda usiopun gu mizeti zote.

VIPIMO VYA KUTUMIA

Mahindi yaliyo kwenye mabunzi Ni vigumu kuyakanga mahindi yaliyo kwenye kukabiliana na wadudu yanapatikana kwa mahindi yaliyopukuswa.

Nafaka (mahindi, ngano na maharagwe) kwenye nafaka iliyopukuswa (mahindi, ngano) changanya gramu 50 (jaza kipimo ulichopewa), kwa kila gramu moja ya nafaka ulichanganya na kifaa cha koleo. Mwingine ACTELIC SUPER nyingine kwenye sehemu ya nje ya gunia lenye nafaka ili kupunguza uwezekano wa shambulizi mpya.

Hifadhi

Dawa ilihadiwe katika viwekeo vyake sili vilivyolingwa, mahali pakavu kuvulini mbali na wadudu, vyakula na mifugo.

WEEKI MBALUNA WATOTO NA WATU

Dawa imalizi kopo: Zika kiasi kidogo cha dawa iliyobaki ndani ya masitomo mbali mbali katika sehemu zisizo na mimea au maji. Safisha mitebe na mapipa ya dawa viziti kabisa ukimwaga maji yenye dawa ndani ya bomba la kunyunyaza. Ongeza maji mengi kwenye baki halafu umwage ndani mbali na mimea. Safisha viziti bomba baada ya kazi.

Formulated and distributed in Kenya by: Twiga Chemical Industries Limited P.O. Box 30172, Nairobi, Kenya. Tel: 338333.

Manufacture by: Syngenta Crop Protection AG Basle, Switzerland

Actellic Super Dusting Powder is a registered trade mark of a Syngenta Group Company

Registration No. / Nambari ya kusajili: PCPB(CR) 0150

Actellic Super Dusting Powder is a registered trade mark of a Syngenta Group Company

Formulated and distributed in Kenya by: Twiga Chemical Industries Limited P.O. Box 30172, Nairobi, Kenya. Tel: 338333.

Manufacture by: Syngenta Crop Protection AG Basle, Switzerland

Actellic Super Dusting Powder is a registered trade mark of a Syngenta Group Company

ACTELIC SUPER

(ACTELIC SUPER Dust ni kiuadudu kwa ajili ya kudhibiti wadudu waharibifu kama dhumuzi na mdundutumbo kwenye mazaio ya mahindi yaliyohifa dhiwa ghalani).

GUARANTEE /DHAMANA

Pirimiphos-methyl 16g/kg
Permethrin 3g/kg

REGISTRATION NUMBER/ NAMBARI YA KUSAJILIWA PCPB (CR) 0150 COMMERCIAL AND AGRICULTURAL CLASS (KUNDI LA BIASHARA NA KILIMO

KEEP LOCKED OUT OF REACH OF CHILDREN FUNGIA MBALUNA WATOTO

HOLOGRAM

READ THE ATTACHED LEAFLET BEFORE USING THE PRODUCT

DATE OF MANUFACTURE/ TARIEHI YA KUTENGENZWA: EXPIRY/UMIA KABA LA: BATCH NO./ NAMBARI YA KIFURUSHI

SHELF LIFE: 2 years from the date of manufacture stored in original sealed containers

Manufacturer/Watengenezaji: Syngenta Crop Protection AG Basle, Switzerland

* Registered Trademark of a Syngenta Group Company

In case of an emergency Call Toll free No. 0800 720021 / 0800 730030(24hrs)

Kukiwa na jambo la dharuru piga simu isiotozwa Nambari: 0800 720021 / 0800 730030(masa 24)

General Information

Broad spectrum insecticidal dust for the control of stored maize and small grain cereals including control of larger grain borer.

Mode of Action

Pirimiphos-methyl rapidly penetrates the insect cuticle and disrupts nerve conduction via the inhibition of acetylcholinesterase. Permethrin is considered to act by interfering with the nicotinic acetyl choline receptor of the nervous system of insects.

Spectrum of activity

Actellic Super Dust controls a wide range of key pests such as the larger grain borer, moths, beetles, mites and weevils. It controls existing infestations and gives long-term protection against re-infestation. Actellic Super Dust is effective under tropical, humid conditions. It persists for several months on grain and inert surfaces. Actellic Super Dust can be applied direct to grain, and to bags.

Resistance Management

In Stored Products, Actellic Super Dust is effective against OP-resistant strains (e.g. those resistant to malathion).

DIRECTIONS FOR USE

BEFORE TREATMENT THE GRAIN SHOULD BE DRY PREFERABLY WITH MOISTURE CONTENT OF LESS THAN 14% FOR EFFECTIVE PEST CONTROL WITH ACTELIC SUPER. FOR BEST RESULTS GRAIN SHOULD BE TREATED IMMEDIATELY AFTER SHELLING AT THE TIME OF STORING. Established infestations are difficult to control. Good coverage is important though the fumigant action partially compensates for this. Treated grain is protected for at least six months.

RATES FOR USE

Cob-maize It is not recommended to use ACTELIC SUPER on cob maize to control Larger Grain Borer. Better control is obtained on shelled maize.

Shelled grain (maize, wheat) Mix 50g (fill the provided measure to the mark) for every 50kg bag using a shovel; the grain and ACTELIC SUPER have to be admixed for effective control of the larger grain borer. Sprinkle some ACTELIC SUPER on the outside of the bag to reduce infestation.

Storage conditions

Store away from sun and damp, in a well-ventilated area apart from food and feed, under lock and key. DO NOT contaminate water, food or feed. DO NOT stack containers in piles more than 2 m high, in order to avoid product compaction or damage. Avoid temperatures above 35°C.

Safety Aspects

Precautionary measures KEEP LOCKED UP OUT OF REACH OF CHILDREN and other unauthorised persons, and animals.

Operator safety

ACTELIC SUPER contains pirimiphos-methyl, which is an organophosphate. Do not use the product if a doctor has advised you not to work with organophosphates, carbamate or thiocarbamate compounds. AVOID getting product in contact with skin, eyes or clothing. DO NOT inhale dust. DO NOT eat, drink or smoke during work. Wash hands and face before eating, drinking or smoking.

WEAR protective clothing: overalls, impermeable gloves, and solid footwear.

AFTER WORK, thoroughly clean protective clothing and equipment used. Wash entire body and change clothing. Wash contaminated clothing before re-use.

Environment and wildlife

DO NOT contaminate water used for irrigation or domestic purposes and water areas, such as ponds, ditches, lakes, drainage systems, etc. by disposal of product waste.

Spillage and leakage

Remove heavily contaminated soil layer; dispose of in a safe place away from water supplies. Collect up spill material thoroughly and dispose it in a safe place or put it in a waste bin or in a plastic bag and dispose of in a landfill site approved for pesticides.

Label: Spill product must not be re-used. Product and container disposal Provided there are no local legal or other rules and regulations, the following procedure is recommended. Dispose of surplus product on a landfill site approved

Net Content/Ujazo: 500gms/Gramu 500

syngenta.

WARRANTY

Sellers guarantee is limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use of this product and accepts the product on that basis.

HIKI KIFUNGUO LAZIMA KIWE KWENYE RANGI NYEUPU NA SAMAWATI

ACTELIC SUPER

(ACTELIC SUPER Dust ni kiuadudu kwa ajili ya kudhibiti wadudu waharibifu kama dhumuzi na mdundutumbo kwenye mazaio ya mahindi yaliyohifa dhiwa ghalani).

GUARANTEE /DHAMANA

Pirimiphos-methyl 16g/kg
Permethrin 3g/kg

REGISTRATION NUMBER/ NAMBARI YA KUSAJILIWA PCPB (CR) 0150 COMMERCIAL AND AGRICULTURAL CLASS (KUNDI LA BIASHARA NA KILIMO

KEEP LOCKED OUT OF REACH OF CHILDREN FUNGIA MBALUNA WATOTO

HOLOGRAM

READ THE ATTACHED LEAFLET BEFORE USING THE PRODUCT

DATE OF MANUFACTURE/ TARIEHI YA KUTENGENZWA: EXPIRY/UMIA KABA LA: BATCH NO./ NAMBARI YA KIFURUSHI

SHELF LIFE: 2 years from the date of manufacture stored in original sealed containers

Manufacturer/Watengenezaji: Syngenta Crop Protection AG Basle, Switzerland

* Registered Trademark of a Syngenta Group Company

In case of an emergency Call Toll free No. 0800 720021 / 0800 730030(24hrs)

Kukiwa na jambo la dharuru piga simu isiotozwa Nambari: 0800 720021 / 0800 730030(masa 24)

Net Content/Ujazo: 500gms/Gramu 500

syngenta.

WARRANTY

Sellers guarantee is limited to the terms set out on the label and subject thereto, the buyer assumes the risk to persons or property arising from the use of this product and accepts the product on that basis.

IPRAF LABEL APPROVED SUBJECT TO SUBMISSION OF FINAL LABEL AND PACKAGE VALUATION 3 MONTHS ONLY

04 DEC 2015

PEST CONTROL PRODUCTS BOARD